

[JP,2002-058483,A]

Japanese (PDF)

File Wrapper Information

[Translation done.]

FULL CONTENTS CLAIM + DETAILED DESCRIPTION TECHNICAL FIELD PRIOR ART TECHNICAL PROBLEM MEANS EXAMPLE

[Translation done.]

Disclaimer:

This English translation is produced by machine translation and may contain errors. The JPO, the INPI, and those who drafted this document in the original language are not responsible for the result of the translation.

Notes:

1. Untranslatable words are replaced with asterisks (****).
2. Texts in the figures are not translated and shown as it is.

Translated: 13:11:09 JST 11/09/2008

Dictionary: Last updated 10/08/2008 / Priority: 1. Biotechnology / 2. Medical/Pharmaceutical sciences / 3.

Technical term

EXAMPLE

[Example] Hereafter, the example of an examination is given and this invention is explained in detail.

[0041]

[The example 1 of an examination] The real-time step RT-PCR method adopted by the creation exam of the calibration curve by the real-time detecting method carries out simultaneously the reaction from which a maximum of 96 differed using 96 wells, and can measure a maximum of 96 specimen at once. Moreover, RT reaction and a PCR reaction can be performed within 1 tube.

[0042] The principle of this quantum uses FRET produced when a wavelength field overlaps between the fluorescence wavelength of one pigment (reporter pigment), and the excitation light wavelength of the pigment (quencher pigment) of another side using two kinds of adjoining fluorochromes. Hybridization is carried out to cDNA of the specific glutathione-S-transferase origin in which the probe (TaqMan probe) combined with both ends amplified two sorts of fluorochromes which produce FRET by PCR. The extension reaction of PCR starts in this status, and it is Taq. A TaqMan probe is hydrolyzed by 5'-3' endonuclease activity which a DNA polymerase has. Since a reporter pigment ****s, the physical distance between quencher pigments arises, the fluorescence intensity of the reporter pigment controlled by FRET increases and the increase in this fluorescence intensity is proportional to the increase of stock of the amplification product of PCR By measuring this for every PCR reaction, a desired quantum becomes possible.

[0043] In the exam, FAM was used as a reporter pigment and TAMRA was used for the three-dash terminal side as a quencher pigment at the five prime end side which is a probe. This etc. performed combination of each pigment, and creation of the TaqMan probe by this according to the procedure (Genome Res., 6 (10), and 986 (1996)) given in literature.

[0044] Moreover, the oligonucleotide as each primer and a probe was compounded using the substrate (dNTP) and the regular reagent using DNA / RNA-biosynthesis machine made from ABI as an automatic synthesizer.

[0045] As a specimen RNA, all the RNA refined from adult's brain, adult's kidney, adult's lungs, adult's small intestine, embryonic liver, and the adult's liver Poole was used. In addition, each of all the RNA, such as this, was purchased from Clonetec (Clontech

Laboratories, Inc.).

[0046] All the RNA refined from adult's brain, adult's kidney, adult's lungs, adult's small intestine, embryonic liver, and the adult's liver Poole was diluted with a RNase freelancer's water, and was made into 20microg/mL. Then, an equivalent amount of these six kinds were mixed every, and it carried out to calibration curve creation. Henceforth, it diluted with the common ratio 5 times using the yeast tRNA of 50microg/mL (Yeast tRNA, product made by GIBCO). 5microL was used for measurement.

[0047] RT-PCR reactions are a 300nM forward primer, a 900nM reverse primer, and a 200nM TaqMan probe. Using included TaqMan One-Step RT-PCR Master Mix Reagents Kit (PE Applied Biosystems), [the system of 50microL / tube] ABI PRISM™7700 It carried out in Sequence Detection System (PE Applied Biosystems).

[0048] After it kept it warm in 30 minutes at 48 degrees C and temperature conditions kept them warm in 10 minutes at 95 degrees C, they were performed for 15 seconds at 95 degrees C, performed the cycle for 1 minute 50 times at 60 degrees C, and measured fluorescence intensity for every cycle.

[0049] The result (calibration curve) of the above-mentioned examination done using the primer pair of arrangement and probe which are shown in each arrangement number to each above mentioned enzyme is shown in Table 1.

[0050]

[Table 1]

酵素	検量線		相関係数 (反応液量: 50 μ L)	定量限界 (pg 全 RNA)
	傾き	切片		
(1) GSTP1	-3.40	38.20	1.00	1.28
(2) GSTT1	-3.21	37.78	1.00	6.4
(3) GSTT2	-4.01	43.15	0.99	32
(4) GSTM1b	-3.80	45.61	0.99	32
(5) GSTM2	-3.84	45.19	1.00	160
(6) GSTM3	-3.31	39.59	1.00	32
(7) GSTM4	-3.55	39.63	1.00	32
(8) GSTM5	-4.31	47.95	1.00	800
(9) GSTA1-1	-3.68	42.52	1.00	4000
(10) GSTA2	-4.65	46.29	1.00	800
(11) GSTA3	-6.38	62.04	0.99	4000
(12) GSTA4	-3.31	39.94	1.00	6.4
(13) MGST1	-3.29	36.29	1.00	6.4
(14) MGST2	-3.66	41.41	1.00	32
(15) MGST3	-3.33	37.03	1.00	32
(16) MGST111	-2.88	41.54	1.00	32
(17) GSTZ 1	-3.27	38.95	1.00	6.4

[0051] When the calibration curve diluted with the common ratio 5 times from 100000pg all RNA / the amount of 50microL reaction mixture was created from Table 1, it is about GSTP1, 1. It had a fixed quantity of sexes to 28pg all RNA / the amount of 50microL reaction mixture, and the coefficient of correlation (r) of the calibration curve which made the quantitation limit 4000pg all RNA from 1.28pg about other enzyme was 0.99 or more.

[0052]

[Layout Table] SEQUENCE LISTING <110> Ostuka Pharmaceutical Factory Inc. <120> The measuring method of glutathione-S-transferase, A probe and a kit <130> for that 00P1072 <160> 51<210> 1 <211> 26<212> DNA <213> human GSTP1 gene <400>

1aggacctccg ctgcaaatac atctcc 26<210> 2 <211> 26<212> DNA <213> human GSTT1 gene <400> 2 ttgcctcgagg acaaggcttcc ccagaa 26<210> 3<211> 26< DNA[212>] <213> human GSTT2 gene <400> 3tttagagetgc gcaccgtgga ttgg 26<210> 4 <211> 31<212> DNA <213> human GSTM1B gene <400> 4 accatggaca accatatgca gctgggcatg a 31<210> 5<211> 29<212> DNA<213> human GSTM2 gene<400> 5tttatggaca gccgtatgca gctggccaa 29<210> 6<211> 31<212> DNA<213> huma n GSTM3 gene<400> 6taatggattt ccgcacacaa ctgataaggc t 31<210> 7 <211> 33<212> DNA <213> human GSTM4 gene <400> 7tccaaatcgc tggccagagt ctgctacage cct 33<210> 8 <211> 33<212> DNA <213> human GSTM5 gene <400> 8cagtcgtacc agctccatgt ggttatccat aac 33<210> 9 <211> 31<212> DNA <213> human GSTA1-1 gene <400> 9gtatgtccac ctgaggaaaa agatgccaag c 31<210> 10 <211> 29<212> DNA <213> human GSTA2 gene<400> 10tactcaacct gaggacaag atgccaagc 29<210> 11<211> 32<212> DNA<213> human GSTA3 gene<400> 11agaaaaacaaa aagtcgetat t tccctgcct tc 32<210> 12 <211> 29<212> DNA <213> human GSTA4 gene <400> 12 tggcaaggaaac ctcaggaga gaacctgta 29<210> 13<211> 30< DNA[212>] <213> human MGST1 gene <400> 13acagccatcc tgcacttcag actatttgc 30<210> 14 <211> 27<212> DNA <213> human MGST2 gene <400> 14ctcgccctgt cagcaaaatg attttgc 27<210> 15 <211> 26<212> DNA <213> human MGST3 gene <400> 15 cacatcttca actgcattca gcgagc 26<210> 16<211> 29<212> DNA<213> human MGST1L1 gene<400> 16tttaggaccca gaaaggagta gacgaagcc 29<210> 17<211> 26<212> DNA<213> human GSTZ 1 gene<400> 17catggagagt tcgaattgct ctggcc 26<210> 18 <211> 21<212> DNA <213> human GSTP1 gene <400> 18 ctgggtggaca tggtaatga c 21<210> 19 <211> 23<212> DNA <213> human GSTP1 gene <400> 19cgcctcatag ttgggtgtaga tga 23<210> 20 <211> 21<212> DNA <213> human GSTT1 gene <400> 20 agagttggat gtgaccctgc a 21<210> 21 <211> 23<212> DNA <213> human GSTT1 gene <400> 21tcagctaagg agatgtgagg acc 23<210> 22<211> 19<212> DNA<213> human GSTT2 gene<400> 22ccaagaagaa tggcattccc 19<210> 23<211> 19<212> DNA<213> human GSTT2 gene<400> 23ttgcctttgt gctgccctt 19<210> 24 <211> 21<212> DNA <213> human GSTM1B gene <400> 24agcaacgcca tcttgcata c 21<210> 25 <211> 22<212> DNA <213> human GSTM1B gene <400> 25 tacgggtggag gtcaaggaca tc 22<210> 26 <211> 21<212> DNA <213> human GSTM2 gene <400> 26aatgcctatcc tgccgtatcat t 21<210> 27 <211> 22<212> DNA <213> human GSTM2 gene <400> 27 tgcctccccaa gaaactgtga gt 22<210> 28 <211> 23<212> DNA<213> human GSTM3 gene<400> 28cgagttggaca tcatagagaa cca 23<210> 29<211> 23<212> DNA<213> human GSTM3 gene<400> 29agc tcttcca agtactgagg ctt 23<210> 30 <211> 21<212> DNA <213> human GSTM4 gene <400> 30ttggagaacc aggtatggaa c 21<210> 31 <211> 22<212> DNA <213> human GSTM4 gene <400> 31 tccccagga actgtgagaa gt 22<210> 32 <211> 21<212> DNA <213> human GSTM5 gene <400> 32ttcccaatct gcccacttgc a 21<210> 33 <211> 22<212> DNA <213> human GSTM5 gene <400> 33 cctccaagta tttgggttc ag 22<210> 34 <211> 20<212> DNA <213> human GSTA1-1 gene<400> 34atgtccagc aagtgcctat 20<210> 35<211> 19<212> DNA<213> human GSTA1-1 gene<400> 35actggagtca agctctcg 19<210> 36<211> 22< DNA[212>] <213> human GSTA2 gene <400> 36cagaccagag ccattctcaa ct 22<210> 37 <211> 23<212> DNA <213> human GSTA2 gene <400> 37 aaggctagag tcaagcttt cca 23<210> 38<211> 22< DNA[212>] <213> human GSTA3 gene <400> 38agatgccaag attgccttga tc 22<210> 39 <211> 25<212> DNA <213> human GSTA3 gene <400> 39 ctctttctaa agcttttgc tctgc 25<210> 40<211> 21< DNA[212>] <213> human GSTA4 gene <400> 40 agttggtaca gacccgaagc a 21< 210> 41<211> 22<212> DNA <213> human GSTA4 gene <400> " *** 1agtccagca gatccag tgt cc 22<210> 42<211> 21< DNA[212>] <213> human MGST1 gene <400> 42tattcccttga gtggcccttgc a 21<210> 43 <211> 22<212> DNA <213> human MGST1 gene <400> 43 aatgggtgtgg tagatccgtg ct 22<210> 44 <211> 22<212> DNA <213> human MGST2 gene <400> 44ctgcgtggct ctgtctctat tc 22<210> 45 <211> 22<212> DNA <213> human MGST2 gene <400> 45 ttgtgtgcc cggaaatactc tc 22<210> 46 <211> 21<212> DNA <213> human MGST3 gene<400> 46tacagcaacgg accctgaaaaa t 21<210> 47<211> 21<212> DNA <213> human MGST3 gene<400> 47acttccaacg tttctggtg g 21< 210> 48<211> 21<212> DNA <213> human MGST1L1 gene <400> 48ggaacgacat ggagaccatc t 21<210> 49 <211> 19<212> DNA <213> human MGST1L1 gene <400> 49 agtgcattcca ggcgacaaa 19<210> 50 <211> 21<212> DNA <213> human GSTZ1 gene<400> 50tctattcc gaagtcctg c 21<210> 51<211> 22<212> DNA<213> human GSTZ1 gene<400> 51ttcagtgcctt ggaagtcctt ag 22

[Translation done.]

[Report Mistranslation](#)

[Japanese \(whole document in PDF\)](#)